

## Supplementary Materials for

# Incommensurate spin correlations in highly oxidized cobaltates $\text{La}_{2-x}\text{Sr}_x\text{CoO}_4$

Z. W. Li<sup>1</sup>, Y. Drees<sup>1</sup>, C. Y. Kuo<sup>1</sup>, H. Guo<sup>1</sup>, A. Ricci<sup>2</sup>, D. Lamago<sup>3,4</sup>, O. Sobolev<sup>5,6</sup>, U. Rütt<sup>2</sup>, O. Gutowski<sup>2</sup>, T. W. Pi<sup>7</sup>, A. Piovano<sup>8</sup>, W. Schmidt<sup>8,9</sup>, K. Mogare<sup>1</sup>, Z. Hu<sup>1</sup>, L. H. Tjeng<sup>1</sup>, and A. C. Komarek<sup>1,\*</sup>

<sup>1</sup> Max-Planck-Institute for Chemical Physics of Solids, Nöthnitzer Str. 40, 01187 Dresden, Germany

<sup>2</sup> Deutsches Elektronen-Synchrotron DESY, Notkestr. 85, 22603 Hamburg, Germany

<sup>3</sup> Forschungszentrum Karlsruhe, Institut für Festkörperphysik, P.O.B. 3640, D-76021 Karlsruhe, Germany

<sup>4</sup> Laboratoire Léon Brillouin, CEA/CNRS, F-91191 Gif-sur Yvette Cedex, France

<sup>5</sup> Forschungsneutronenquelle Heinz Maier-Leibnitz (FRM-II), TU München, Lichtenbergstr. 1, D-85747 Garching, Germany


<sup>6</sup> Georg-August-Universität Göttingen, Institut für Physikalische Chemie, Tammannstrasse 6, D-37077 Göttingen, Germany

<sup>7</sup> National Synchrotron Radiation Research Center (NSRRC), Hsinchu 30077, Taiwan

<sup>8</sup> Institut Laue-Langevin (ILL), 71 avenue des Martyrs, F-38042 Grenoble Cedex 9, France

<sup>9</sup> Jülich Centre for Neutron Science JCNS, Forschungszentrum Jülich GmbH, Outstation at ILL, 71 avenue des Martyrs, F-38042 Grenoble Cedex 9, France


\* Alexander.Komarek@cpfs.mpg.de



**Fig. S1** The Co- $L_{2,3}$  XAS spectrum of  $\text{La}_{1.2}\text{Sr}_{0.8}\text{CoO}_4$  is shown together with XAS spectra of CoO,  $\text{NdCaCoO}_4$  and  $\text{Sr}_2\text{Co}_{1/2}\text{Ir}_{1/2}\text{O}_4$  reference samples. The pre-peak at 778 eV can be attributed to the presence of a  $\text{Co}^{2+}$  species. The experimental Co- $L_{2,3}$  spectrum of  $\text{La}_{1.2}\text{Sr}_{0.8}\text{CoO}_4$  can be decomposed into 77%  $\text{Co}^{3+}$  content, denoted as ‘difference’ (black circles) and 23%  $\text{Co}^{2+}$  contribution (cyan line). These measurements show that our  $\text{La}_{1.2}\text{Sr}_{0.8}\text{CoO}_{4+\delta}$  sample has only a tiny oxygen deficiency ( $\delta = -0.015 \pm 0.005$ ). After subtraction of the  $\text{Co}^{2+}$  contribution (denoted as ‘difference’), we were able to estimate the HS content for the  $\text{Co}^{3+}$  species. Therefore, we used  $\text{NdCaCoO}_4$  and  $\text{Sr}_2\text{Co}_{0.5}\text{Ir}_{0.5}\text{O}_4$  [S1,S2] as a  $\text{Co}^{3+}$  LS and  $\text{Co}^{3+}$  HS reference material, respectively. The lower part of this figure shows that the difference spectrum can be nicely reproduced by the weighted sum (magenta) of the  $\text{Co}^{3+}$  HS spectrum (green line) and the  $\text{Co}^{3+}$  LS spectrum (blue line). Thus, the presence of ~19%  $\text{Co}^{3+}$  ions in the HS state and ~58%  $\text{Co}^{3+}$  ions in the LS state is revealed by this modelling of the ‘difference’ spectrum (besides 23%  $\text{Co}^{2+}$  ions).

## References

- [S1] X. Ou and Hua Wu, Phys. Rev. B **89**, 035138 (2014)
- [S2] unpublished results



**Fig. S2** Fourier transformation of charge ordering pattern used for numerical simulations in Fig. 6 (a). The expected intensity for charge correlations is shown in a logarithmic colour-contour plot. Since only the modulations of the Co-oxygen bond distances can be seen with neutrons and not the charges itself, the values are arbitrary.

## Supplementary Video Legends

### Supplementary Video #1: **Low energy magnetic excitations**

The movie shows a simulation of spin excitations at  $\sim 1.3$  meV within our nano phase separation model for  $\text{La}_{1.3}\text{Sr}_{0.7}\text{CoO}_4$ .

### Supplementary Video #2: **Higher energy magnetic excitations**

The movie shows a simulation of spin excitations at  $\sim 8.4$  meV within our nano phase separation model for  $\text{La}_{1.3}\text{Sr}_{0.7}\text{CoO}_4$ .